

Comparing maternal pre-pregnancy weight classification methods and the effect on gestational weight gain classification among MI adolescents, 2003-2007

Patricia McKane, DVM MPH
Epidemiologist –WIC & Title X Family
Planning Programs
Michigan Department of Community Health

Mckanep@michigan.gov



Acknowledgements

■ Co-investigators

■ Violanda Grigorescu, MD, MSPH¹

■ Kobra Eghtedary, PhD²

1 Division of Genomics, Perinatal Health and Chronic Disease
Epidemiology, Michigan Department of Community Health

2 WIC Division, Bureau of Family, Maternal and Child Health,
Michigan Department of Community Health

■ Study was granted exemption from the Michigan
Department of Community Health IRB

Outline

- Introduction
- Methods
- Results
- Discussion
- Public Health Significance

3

Introduction

- Pregnancy weight gain is an important determinate of infant & maternal outcomes¹⁻³
- 1990-Institute of Medicine (IOM) developed pregnancy weight gain recommendations based on adult Body Mass Index (BMI) categories¹
- Discordance between IOM 1990 and CDC BMI weight for age identified^{5,6}
- 2007-IOM Workshop expressed concern that adult BMI categories were inappropriate for adolescents⁴
- 2009 -IOM revised pregnancy weight gain recommendations and suggest that adolescents continue to be classified by adult BMI categories⁷

4

Introduction

Comparison of Institute of Medicine pre-pregnant adult BMI categories and CDC BMI percentile categories (age/gender specific)

Pre-pregnancy BMI	IOM 1990	IOM 2009	CDC BMI percentile
Underweight	< 19.8	< 18.5	< 5th
Normal weight	19.8 - 26.0	18.5 – 24.9	5 th - < 85th
Overweight	26.1 - 29.0	25.0 - 29.9	85 th - < 95th
Obese	≥ 29.1	≥ 30.0	≥ 95th

5

Introduction

Institute of Medicine pregnancy weight gain recommendations based on pre-pregnancy BMI

Pre-Pregnancy BMI	IOM 1990 rec. wt. gain (lbs)	IOM 2009 rec. wt. gain (lbs)
Underweight	28 - 40	28 - 40
Normal weight	25 – 35	25 - 35
Overweight	15 – 25	15 -25
Obese	At least 15	11-20

Note that the IOM 1990 did not establish an upper limit for obese women; however, the upper limit of 25 pounds was used in PNSS data analysis.


6

Study Questions

- Is there discordance between BMI categorization methods among pre-pregnant adolescents?
- Is discordance associated with inappropriate weight gain based on IOM 1990 and/or 2009 recommendations?

7

Methods



8

Methods

- Pre-Pregnancy BMI (kg/m²) based on:
 - Pre-Pregnancy weight (kg)
 - Self-reported
 - Correlated with prenatal weight (measured at WIC enrollment)
 - Pearson's correlation coefficient ranged from 0.919 to 0.927 (per year)
 - Mean difference \pm standard error
 - Overall ranged from 5.0 ± 0.6 to 5.5 ± 0.12 kg
 - Increased by WIC Trimester 2.0 ± 0.7 kg -1st trimester
 4.4 ± 0.9 kg-2nd trimester & 11.3 ± 0.15 -3rd trimester

9

Methods-outcomes

- Discordance
 - BMI CDC vs. BMI 90 or BMI 09
 - Also predictor variable for 2nd study question
- Weight gain categorized as less than recommended, recommended or more than recommended based on:
 - IOM 90
 - IOM 09

10

Methods-independent variables

- Age
- Race/Ethnicity
- Previous pregnancy
- Pre-Pregnancy weight
 - BMI 90 –categorized based on IOM 1990 criteria
 - BMI 09 –categorized based on IOM 2009 criteria
 - BMI CDC –categorized base on the CDC age/gender specific BMI tables
- Poverty Level
- Marital Status

SAS 9.1 (SAS Institute Inc., Cary, NC) was used for statistical analyses

11


Results- Distribution of selected characteristics

Characteristics	N	%
Age (years)		
10-15	3552	5.6
16-17	14637	23.0
18-19	29500	46.6
20	15720	25.0
Race/Ethnicity		
White, non-Hispanic	35900	56.6
Black, non-Hispanic	19888	31.4
Hispanic	5887	9.3
Asian/Pacific Islander	452	0.7
American Indian/Alaska Native	275	0.4
Multi-racial, non-Hispanic	1007	1.6
All	63409	

12

Results


Prevalence of Pre-pregnancy BMI among adolescents ages 10-20 by CDC age/gender specific categories, IOM 1990 categories and IOM 2009 categories, MI PNSS 2003-2007



13


Results-Discordance

Prevalence of discordance of BMI by selected characteristics among adolescents 10-20 years of age, MI PNSS 2003-2007



Results-Logistic Regression

Adjusted and crude effects of characteristics on BMI discordance (IOM 1990)



•Adjusted for age, race/ethnicity, previous pregnancy Passed Hosmer and Lemeshow test, but c value = 0.547 Reference: White, non-Hispanic, Age 20 years

15

Results-Logistic Regression

IOM 2009 OR;95% CI†

■ Maternal Age


- 10-15 years 1.155; 1.035, 1.287
- 16-17 years 0.449; 0.413, 0.488
- 18-19 years 0.749;0.704, 0.797

†Model includes age- all other effects eliminated by backward selection. Compared to 20 years of age

16


Results-Weight gain IOM 1990

Distribution of selected characteristics and weight gain based on IOM 1990 among adolescents 10-20 years of age, MI PNSS 2003-2007



Results – Multinomial Regression IOM 1990


Estimated effects of demographic characteristics and BMI discordance on weight gain categorized as **less** than recommended, adolescents ages 10-20 years MI PNSS 2003-2007



Model includes maternal age, race/ethnicity, BMI (CDC), marital status, previous pregnancy and BMI discordance between IOM 1990 and CDC BMI. OR = 1.0 for: No BMI Discordance, Normal weight, White, non-Hispanic, and 20 years.

Results – Multinomial Regression IOM 1990

Estimated effects of demographic characteristics and BMI discordance on weight gain categorized as **more** than recommended, adolescents ages 10-20 years MI PNSS 2003-2007




Model includes maternal age, race/ethnicity, BMI (CDC), marital status, previous pregnancy and BMI discordance between IOM 1990 and CDC BMI OR=1.0 for: No BMI Discordance, No previous pregnancy, single Normal weight, White, non-Hispanic

19


Results-Weight gain IOM 2009

Distribution of selected characteristics and weight gain based on IOM 2009 among adolescents 10-20 years of age, MI PNSS 2003-2007



Results – Multinomial Regression IOM 2009

Estimated effects of demographic characteristics and BMI discordance on weight gain categorized as **less** than recommended, adolescents ages 10-20 years MI PNSS 2003-2007




Model includes maternal age, race/ethnicity, BMI (CDC), marital status, previous pregnancy, income and BMI discordance between IOM 2009 and CDC BMI. OR =1.0 for: No BMI Discordance, Normal weight, White, non-Hispanic, and 20 years.

21

Results – Multinomial Regression IOM 2009

Estimated effects of demographic characteristics and BMI discordance on weight gain categorized as **more** than recommended, adolescents ages 10-20 years, MI PNSS 2003-2007



Model includes maternal age, race/ethnicity, BMI (CDC), marital status, previous pregnancy, income and BMI discordance between IOM 2009 and CDC BMI. OR =1.0 for: Poverty level > 100%, single Normal weight, White, non-Hispanic and Age 20 years.

22

Conclusion

- Discordance between the three methods of BMI classification was found
 - Less for IOM 2009
- Discordance was associated with inappropriate weight gain based IOM recommendations
- Racial disparities were evident
- Underweight adolescents at increased odds of less than recommended weight gain

23

Public Health Implications

- BMI categorization based on the CDC age/gender specific percentiles for pregnant adolescents may be more appropriate.
- Gestational weight gain recommendations specifically for adolescents must be further explored and thus better understood.

24

References

1. Institute of Medicine. *Nutrition During Pregnancy*. Washington DC: National Academies Press; 1990. Accessed online April 7, 2010 at http://books.nap.edu/openbook.php?record_id=1451&page=R1
2. Viswanathan M, Siega-Riz AM, Moos MK, Deierlein A, Mumford S, Knaack J, Thieda P, Lux LJ, Lohr KN. Outcomes of maternal weight gain. Evidence Report/Technology Assessment. 2008; 168:1-223.
3. Oken E, Taversas EM, Kleinman KP, et al. Gestational weight gain and child adiposity at age 3 years. *Am J Obstet Gynecol*. 2007; 196(4):322.1-8
4. National Research Council and Institute of Medicine. Influence of pregnancy weight on maternal and child health. Workshop report. Washington DC: National Academies Press. 2007. Accessed online April 7, 2010 at http://books.nap.edu/openbook.php?record_id=11817
5. Fernandez ID, Olson CM & De Ver Dye, T. Discordance in the assessment of prepregnancy weight status of adolescents: A comparison between the Centers for Disease Control and Prevention sex- and age-specific body mass index classification and the Institute of Medicine-based classification used for maternal weight gain guidelines. *J Am Diet Assoc*. 2008; 108(6): 998-1002
6. Groh S. Are the Institute of Medicine recommendations for gestational weight gain appropriate for adolescents? *JOGNN*. 2007; 36: 21-27
7. Committee to reexamine IOM pregnancy weight guidelines. *Weight Gain During Pregnancy: Reexamining the Guidelines* Washington DC: National Academies Press; 1990. Accessed online April 7, 2010 at <http://www.iom.edu/Reports/2009/Weight-Gain-During-Pregnancy-Reexamining-the-Guidelines.aspx>

25

Thank you

26